

Prontosan® Wound Gel | Prontosan® Wound Gel X

Cleansing and moistening of skin wounds and burns. For the prevention of biofilm.



- Removes & prevents biofilm
- Prevents infections
- Reduces healing time
- Painless dressing changes
- Well-known substances with low allergenic potential
- Compatible with commonly used wound dressings
- Can be used up to 8 weeks after first opening


While necrotic tissue and slough can be optimally removed by Prontosan® Wound Irrigation Solution, it is necessary to first remove heavy encrustations by keeping the wound moist for a longer period of time. For this task, Prontosan® Wound Gel and Gel X provide the optimal solution.

When to use which gel

Prontosan® Wound Gel	Prontosan® Wound Gel X
30 ml pod	250 g tube
Fluid	Highly viscous
For smaller, deep and tunneling wounds, as well as for difficult to access wounds	For large surface area wounds, e.g. <ul style="list-style-type: none"> ➤ Ulcus cruris ➤ Decubitus
For burn wounds I. & II. degree	For all burn wounds, incl. III. & IV. degree
If little quantity is required	When large quantities are required Where Prontosan® Wound Gel is too fluid

HINTS AND TIPS

All wounds should in principle first be rinsed and cleansed with Prontosan® Wound Irrigation Solution, Prontosan® Gel and Gel X remain on the wound until the next dressing change. It therefore has a long acting effect.



Indications

Cleansing, moistening and decontamination of skin wounds and burns

- › Traumatic wounds
- › Postoperative wounds
- › Chronic skin ulcers (e.g. venous, diabetic or pressure ulcers)
- › Thermal wounds
- › Chemical wounds (acid- and alkali-induced)
- › Radiation-induced wounds

Moistening of bandages and wound dressings

Application

The choice between Prontosan® Wound Gel and Gel X allows an optimal application in large surface area as well as deep wounds.

- ① For large surface area wounds apply a 3-4 mm thick layer of the **Prontosan® Wound Gel X** and cover with a secondary dressing.
- ② For the application in deep or tunneling wounds, wound cavities and difficult to access areas, apply a 3-5 mm layer of **Prontosan® Wound Gel** and cover with a secondary dressing.

Prontosan® Wound Gel



	Size	Pcs/Pack	Article number
Prontosan® Wound Gel	30 ml	1	400505
Prontosan® Wound Gel X	50 g	1	400517
Prontosan® Wound Gel X	250 g	1	400508

Composition: 0.1 % Polyaminopropyl Biguanide (Polihexanide), Betaine Surfactant, Purified Water, Glycerol, Hydroxyethylcellulose