

TOXI-PLUS LIQUID

Composition

Each litre contains:

Poly [1-(2-oxo-1-pyrroli-dinyl)ethyle] 3.0%, Bis [(sonicotinal-doximin) 1-methlye] 0.3%

Indication


As a Toxin Binder for mycotoxin e.g. Aflatoxin B, B2, G1, G2, Zearalenone, T-2 etc that will lead to mortality and poor production of the animals.

Act as Feed Enhancer to utilize feed effectively which contains certain organic acids that will improve the digestibility and absorption of the feed. Better Feed Conversion Ratio and improve body weight gain after recovery from the problems due to fungals, molds and mycotoxins. As an Appetizer for efficient feed intake.

As part of a treatment program, **TOXI-PLUS LIQUID** increases production parameters. Given to broilers during the 7 days, it reduces mortality, increase growth and improves response to vaccination. It results to better carcass quality and fewer losses during transfer when given during the last 7 days before harvest.

Broad Spectrum of Activity

- Reduces the negative effects of mycotoxins
- Reduces the in-vivo bioavailability of mycotoxins for gastrointestinal absorbance
- Improves animal immune system
- Improves overall animal performance
- Reduces mortality due to mycotoxins
- Improves immunity and response to vaccination
- Better functioning of body organs, especially liver, kidneys and ovaries
- Cure oral lesions caused by fusarial toxins found in poultry
- Reduces the incidence of leg problems
- Improves amino acid absorption



Dosage & Application

Add 100ml of Toxi-Plus Liquid per 1000 litres of drinking water.

Add 20ml of Toxi-Plus Liquid per 200 litres for 7-10 days or until mold and fungal infections have ceased.

Packing

1 L/btl