

N-FLOK 200

controlled medicine

Composition

Each litre contains:
Norfloxacin BP 200gm

Pharmacodynamic

Norfloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E.coli, Haemophilus, Mycoplasma, Pasteurella, and Salmonella. Norfloxacin acts on the DNA-gyrase and interferes with the copy transcription, and recombination of bacteria DNA, thereby leading to the abnormal growth of bacteria and eventually cause to death.

Indication

To be used in infections caused by microorganism sensitive to Norfloxacin in poultry and swine.

Contraindication

Contraindication administrates norfloxacin to animal hypersensitive to it and animal with a seriously impaired liver and / or renal function.

Drug Interaction

Norfloxacin should not be used together with Rifampicin and drugs that's containing Al, Mg, Iron and Calcium Ion. Contraindication concurrent administration with tetracyclines, chloramphenicol, macrolides and lincosamides.

Side Effects / Adverse Reaction

Hypersensitive reactions, Administration to juvenile animals can lead to arthropathy.

Dosage & Application

Carefully mix with drinking water

For Poultry & Breeders

0.5ml / litre daily for 3 days (Continue treatment with 0.25ml / litre daily for 3 days)

Swine & Suckling Swine

0.4ml / 5kg bodyweight daily for 5 days.

Withdrawal Period

Meat: Poultry : 12 days
Swine : 8 days

Packaging

1 L/btl

